



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

health policy makers or payers, and could further cloud the diagnostic certainty of sarcoidosis, which is already a challenge in many situations. The introduction of the term parasarcoidosis would also affect sarcoidosis research by further obfuscating the goals of treatment. Should we research sarcoidosis and parasarcoidosis separately, or should they be considered as one disease? The implication of parasarcoidosis as a separate entity might undermine what little research investment is made in sarcoidosis. Moreover, individuals with sarcoidosis often have to face and struggle with the fact that their disease is not taken seriously.¹⁹ If we, as professionals, speak about parasarcoidosis, this could send the wrong message that we do not believe these symptoms to be part of their disease. A complex condition such as sarcoidosis justifies a holistic approach in which the patient is taken seriously. In idiopathic pulmonary fibrosis, and other fibrotic lung diseases, fatigue has also been acknowledged as a substantial burden.¹⁰ Will this fatigue be called parafibrosis? We should be careful about stigmatising symptoms in this way.

In summary, patients with sarcoidosis tend to be affected by specific organ-related symptoms with functional impairments and by less specific symptoms. Although some disease manifestations are hard to verify, it does not mean that they are not associated with or do not belong to sarcoidosis. What we need is a guideline to assess, quantify, and manage disability in sarcoidosis, and not a new confusing term to label what we, to date, do not completely understand. Because of its complexity, the care of sarcoidosis needs to be personalised and a multidisciplinary approach is recommended.

FB reports consulting fees and speaker honoraria from Boehringer Ingelheim, Bristol Myers Squibb, F Hoffmann– La Roche, Galapagos, and Savara Pharmaceuticals outside of the submitted work. All other authors declare no competing interests.

**Marjolein Drent, Ulrich Costabel, Elliott D Crouser, Johan Grunewald, Francesco Bonella*
m.drent@ildcare.nl

Interstitial Lung Diseases Center of Excellence, Dept of Pulmonology, St Antonius Hospital, Nieuwegein, Netherlands (MD); Department of Pharmacology and Toxicology, Faculty of Health, Medicine and Life Sciences, Maastricht University, Maastricht, Netherlands (MD); ild care foundation research team, 6711 NR Ede, Netherlands (MD); Center for Interstitial and Rare Lung Diseases, Pneumology Department, Ruhrlandklinik, University Hospital, University of Essen, Essen, Germany (UC, FB); European Reference Network (ERN)-LUNG, ILD Core Network (UC, FB); Division of Pulmonary and Critical Care and Sleep Medicine, The Ohio State University, Columbus, OH, USA (EDC); Chair Scientific Advisory Board Foundation of Sarcoidosis Research, Chicago, IL, USA (EDC); Respiratory Medicine Division, Department of Medicine Solna, and Center for Molecular Medicine, Karolinska Institutet, Stockholm, Sweden (JG); Respiratory Medicine, Theme Inflammation and Ageing, Karolinska University Hospital, Stockholm, Sweden (JG)

- 1 Drent M, Strookappe B, Hoitsma E, De Vries J. Consequences of sarcoidosis. *Clin Chest Med* 2015; **36**: 727–37.
- 2 Voortman M, Hendriks CMR, Elfferich MDP, et al. The burden of sarcoidosis symptoms from a patient perspective. *Lung* 2019; **197**: 155–61.
- 3 Drent M, Lower EE, De Vries J. Sarcoidosis-associated fatigue. *Eur Respir J* 2012; **40**: 255–63.
- 4 Voortman M, Fritz D, Vogels OJM, et al. Small fiber neuropathy: a disabling and underrecognized syndrome. *Curr Opin Pulm Med* 2017; **23**: 447–57.
- 5 Judson MA. The three tiers of screening for sarcoidosis organ involvement. *Respir Med* 2016; **113**: 42–49.
- 6 Lower EE, Sturdivant M, Baughman RP. Presence of onconeural antibodies in sarcoidosis patients with parasarcoidosis syndrome. *Sarcoidosis Vasc Diffuse Lung Dis* 2019; **36**: 254–60.
- 7 Judson MA. Screening sarcoidosis patients for occult disease. *Semin Respir Crit Care Med* 2020; **41**: 741–57.
- 8 Moor CC, Kahlmann V, Culver DA, Wijsenbeek MS. Comprehensive care for patients with sarcoidosis. *J Clin Med* 2020; **9**: 9.
- 9 Hendriks CMR, Saketkoo LA, Elfferich MDP, De Vries J, Wijnen PAHM, Drent M. Sarcoidosis and work participation: the need to develop a disease-specific core set for assessment of work ability. *Lung* 2019; **197**: 407–13.
- 10 Kölner-Augustson L, Prior TS, Skivild V, Alestrup A, Bendstrup E. Fatigue in idiopathic pulmonary fibrosis measured by the Fatigue Assessment Scale during antifibrotic treatment. *Eur Clin Respir J* 2020; **8**: 1853658.



COVID-19 vaccination in patients with α 1-antitrypsin deficiency

Published Online
June 17, 2021

[https://doi.org/10.1016/S2213-2600\(21\)00271-X](https://doi.org/10.1016/S2213-2600(21)00271-X)

For more details on COVID-19 data see <https://covid19.who.int/>

COVID-19 is continuing its spread across the world, generating a wake of devastating health, economic, and social consequences. The urgency of the situation has simultaneously driven the development of COVID-19 vaccines to an astonishingly fast pace, with more than 2.1 billion doses administered worldwide). Despite the hugely successful vaccination campaign, the efficacy of COVID-19 vaccines in

patients with genetic pulmonary diseases, such as α 1-antitrypsin deficiency (AATD), have not been delineated. Inherited AATD is characterised by low concentrations of functional α 1-antitrypsin in the blood, predisposing individuals to enzymatic tissue injury and inflammation, most notably in the lungs. Given the unclear timeline for an end to the ongoing pandemic, shedding light onto such issues that might

have potentially fatal consequences for the affected individuals is necessary.

The efficacy of COVID-19 vaccines in individuals with AATD might only be inferred from published studies on influenza and pneumococcal vaccination in this population. In an observational study of 939 participants with AATD, influenza vaccination rates were up to 81.6% in patients with AATD, yet this finding did not translate into a decrease in exacerbation rates of chronic obstructive pulmonary disease or decreased health-care use.¹ In another study, the magnitude of the antibody response to pneumococcal vaccination in patients with severe AATD was no less robust than in healthy volunteers.² The immune responses to antigen provocation in individuals with AATD appear to be far more complex, and protection from SARS-CoV-2 might not necessarily be afforded with vaccination alone.

We have hypothesised that individuals with AATD might derive limited benefit from the current COVID-19 vaccines for several reasons. First, even though vaccination has been prioritised to more vulnerable populations (such as people with AATD), individuals with AATD are usually not included in clinical trials (as reported in ClinicalTrials.gov), and thus the effectiveness and adverse event profile of vaccination in this population are unknown. Clinical scientists should include the AATD population in clinical trials of COVID-19 vaccines to better characterise the safety and efficacy for individuals with AATD. Second, Kueppers³ has shown that the increase in trypsin-inhibiting capacity of serum after injection of typhoid vaccine is largely due to the increase of the α 1-antitrypsin concentration. Consequently, AATD inhibits the quantitative response of the α 1-antitrypsin to such a stimulus.³ Emerging evidence has shown a strong correlation between concentrations of circulating α 1-antitrypsin and the induction of trained immunity.⁴ Data from previous studies of severe acute respiratory syndrome, Middle East respiratory syndrome, and other human respiratory viruses allude to a risk of antibody-dependent enhancement associated with SARS-CoV-2 vaccines and antibody-based interventions. Liu and colleagues⁵ have recently reported that the concentrations of enhancing and neutralising antibodies were higher in patients with severe COVID-19 than in patients with non-severe COVID-19. For individuals with AATD, there is a possibility that

Panel: Currently available therapeutic approaches to manage α 1-antitrypsin deficiency during the COVID-19 pandemic

Lifestyle changes

- Avoid crowded places and mass gatherings
- Wear a medical mask in public spaces
- Patients should be counselled and receive assistance in smoking cessation
- Patients should be advised to avoid environmental risk exposures, such as smog, second-hand tobacco smoke, dusts, and fumes
- Maintain a healthy bodyweight and consume a balanced, vitamin-rich, and high-fibre diet

Pharmacotherapy

- Patients should receive general medical therapy for chronic obstructive pulmonary disease, including bronchodilators, steroids, and oxygen therapy, as per local chronic obstructive pulmonary disease guidelines

Vaccination


- Patients should receive the annual flu vaccine and pneumococcal vaccine as per guidelines
- Patients should also receive vaccination against SARS-CoV-2

Augmentation therapy

- Intravenous plasma-purified α 1-proteinase inhibitor should be considered for the treatment of α 1-antitrypsin deficiency-related lung disease

Surgery

- Lung transplantation could be considered for patients with very severe disease refractory to pharmacotherapy



Kredesign/Science Photo Library

the antibody-dependent enhancement effect might be amplified by SARS-CoV-2 infection or vaccination. Third, AATD eases virus spike protein activation by elastase, resulting in the faster spread of the SARS-CoV-2 subtype with spike 614Gly mutation, which is more virulent and results in greater host morbidity.⁶ Fourth, COVID-19 vaccine efficacy might not reach 100%; governmental negligence, socioeconomic inequalities, personal values, and the looming spectre of a SARS-CoV-2 mutation all contribute to suboptimal vaccination rates, lower than that of the flu vaccination. Recent studies are raising concern that current COVID-19 vaccines might not have sufficient efficacy against the new SARS-CoV-2 variants B.1.1.7 and B.1.351.^{7,8} In particular, the SARS-CoV-2 variant B.1.351 might decrease SARS-CoV-2 vaccine-derived neutralisation of SARS-CoV-2 by 6–86 times.⁹ Finally, Pi^*MZ , Pi^*SZ , or unknown AATD genotype have been associated with a greater odds of unhealthy

behaviours, such as not obtaining the pneumococcal or influenza vaccine, adopting sedentary lifestyles, and smoking.¹⁰ Individuals who do not know their genotype might require additional education and intervention to mitigate the risk of SARS-CoV-2 infection. As the advocacy for COVID-19 vaccination in people with AATD continues, studies need to elucidate a proven vaccine correlation with SARS-CoV-2 strains.

In the meantime, the protective roles of α 1-antitrypsin on lung structure and function, on preventing acute lung injury and acute respiratory distress syndrome, and especially on inhibiting SARS-CoV-2 infection renders alpha₁-proteinase inhibitor a promising candidate for COVID-19 treatment in select populations.^{11,12} For patients with AATD, alpha₁-proteinase inhibitor therapy might solve two problems with one single action because it is also the best candidate drug for the treatment of COVID-19. As the ongoing pandemic persists in the foreseeable future, we strongly advocate that public health officials and health-care professionals should encourage the population of people with AATD to adopt protective behaviours, including lifestyle changes, pharmacotherapy, alpha₁-proteinase inhibitor therapy, surgery, and other therapeutic approaches in addition to COVID-19 vaccine uptake (panel). Patient-centric educational messages for patients with AATD that emphasise the severity of COVID-19, particularly the potential long-term negative health sequelae, are needed. We must fight to ensure that all patients with AATD, regardless of race, ethnicity, immigration status, income, and insurance status, have access to

essential medications and timely and high-quality care in this difficult time.

CY is a former postdoctoral fellow at the University Health Network. HZ declares no competing interests.

*Chengliang Yang, Hedi Zhao
chengliang.yang@utoronto.ca

Latner Thoracic Surgery Research Laboratories, Toronto General Hospital Research Institute, University Health Network, Toronto, ON M5G 1L7, Canada (CY); Faculty of Medicine, McGill University, Montreal, QC, Canada (HZ)

- Campos MA, Alazemi S, Zhang G, Sandhaus RA, Wanner A. Influenza vaccination in subjects with alpha1-antitrypsin deficiency. *Chest* 2008; **133**: 49–55.
- Miravittles M, de Gracia J, Rodrigo MJ, et al. Specific antibody response against the 23-valent pneumococcal vaccine in patients with alpha(1)-antitrypsin deficiency with and without bronchiectasis. *Chest* 1999; **116**: 946–52.
- Kueppers F. Genetically determined differences in the response of alpha-antitrypsin levels in human serum to typhoid vaccine. *Humangenetik* 1968; **6**: 207–14.
- Cirovic B, de Bree LCJ, Groh L, et al. BCG vaccination in humans elicits trained immunity via the hematopoietic progenitor compartment. *Cell Host Microbe* 2020; **28**: 322–34.
- Liu Y, Soh WT, Kishikawa J, et al. An infectivity-enhancing site on the SARS-CoV-2 spike protein targeted by antibodies. *Cell* 2021; published online May 24. <https://doi.org/10.1016/j.cell.2021.05.032>.
- Bhattacharyya C, Das C, Ghosh A, et al. SARS-CoV-2 mutation 614G creates an elastase cleavage site enhancing its spread in high AAT-deficient regions. *Infect Genet Evol* 2021; **90**: 104760.
- Wang P, Nair MS, Liu L, et al. Antibody resistance of SARS-CoV-2 variants B.1.351 and B.1.1.7. *Nature* 2021; **593**: 130–35.
- Madhi SA, Baillie V, Cutland CL, et al. Efficacy of the ChAdOx1 nCoV-19 COVID-19 vaccine against the B.1.351 variant. *N Engl J Med* 2021; **384**: 1885–98.
- Abdool Karim SS, de Oliveira T. New SARS-CoV-2 variants—clinical, public health, and vaccine implications. *N Engl J Med* 2021; **384**: 1866–68.
- Holm KE, Mannino DM, Choate R, Sandhaus RA. Genotype is associated with smoking and other key health behaviors among individuals with alpha-1 antitrypsin deficiency-associated lung disease. *Respir Med* 2018; **143**: 48–55.
- Yang C, Keshavjee S, Liu M. Alpha-1 antitrypsin for COVID-19 treatment: dual role in antiviral infection and anti-inflammation. *Front Pharmacol* 2020; **11**: 615398.
- Ritzmann F, Chitrala P, Krüger N, et al. Therapeutic application of alpha-1-antitrypsin in COVID-19. *Am J Respir Crit Care Med* 2021; published online May 7. <https://doi.org/10.1164/rccm.202104-0833LE>.



Ongoing asthma management in children during the COVID-19 pandemic: to step down or not to step down?

Published Online
July 9, 2021
[https://doi.org/10.1016/S2213-2600\(21\)00235-6](https://doi.org/10.1016/S2213-2600(21)00235-6)

A substantial reduction in asthma exacerbations in both children and adults has been seen in many countries worldwide during the COVID-19 pandemic.^{1,2} The cause of this reduction is likely to be multifactorial, but at least partly due to population-level public health measures, such as physical distancing, masking, and hand washing, which reduce broad viral transmission.² This improvement in asthma control poses an interesting clinical dilemma: should clinicians consider tapering

asthma medications in children during the pandemic in the face of good asthma control? Furthermore, if medicines are reduced during the COVID-19 pandemic, should clinicians return to the pre-COVID-19 schedule as respiratory precautions are gradually relaxed?

Before the pandemic, the evidence strongly supported a step-down therapeutic approach in children (aged >5 years) with good asthma control. The Global Initiative for Asthma (GINA) strategy, under usual circumstances,